

Salinity tolerance of *Aegiceras corniculatum* (L.) Blanco from Gujarat coasts of India

Neha Tusharbai Patel & Amar Nath Pandey

Department of Biosciences, Saurashtra University, Rajkot 360 005, India

Resumen

Correspondence

A. N. Pandey

E-mail: anpandey2001@gmail.com

Tel (O): +91-281-2586419

Tel (M): +91-9427495989

Fax (O): +91-281-2577633

Received: 31 August 2009

Accepted: 26 October 2009

Published on-line: 12 November 2009

Efectos de la salinidad del suelo en el crecimiento, estado hídrico y acumulación de nutrientes en semillas de Aegiceras corniculatum (Myrsinaceae).

Se realizaron experimentos en invernadero para valorar los efectos de la salinidad del suelo en la emergencia, crecimiento, estado hídrico contenido de prolina y acumulación mineral en plántulas de *Aegiceras corniculatum* (L.) Blanco (Myrsinaceae). Se añadió NaCl al suelo y se mantuvo la salinidad a 0,3; 3,9; 7,9; 12,1; 16,1; 19,7; 24,0; 28,0; 32,0; 36,0 y 40,0 dSm⁻¹. Esta especie es muy tolerante a la sal durante la fase de germinación. Las concentraciones bajas de sal estimularon significativamente el crecimiento de las plántulas, siendo su crecimiento óptimo a 24,9 dSm⁻¹ aunque salinidades mayores lo inhibieron. Al aumentar la salinidad, se produjeron los siguientes efectos: el potencial hídrico de los tejidos se volvió negativo. La cantidad de prolina aumentó. La concentración de Na en los tejidos aumentó, mientras que disminuyeron el K, Ca, N y P. También se discute sobre los cambios en los patrones de acumulación de otros nutrientes y los posibles mecanismos para evitar la toxicidad del Na, en respuesta a la salinidad.

Palabras clave: *Aegiceras corniculatum*; Acumulación mineral; Contenido de prolina; Germinación de propágulos; Crecimiento de plántulas; Estado hídrico.

Abstract

Greenhouse experiments were conducted to assess the effects of soil salinity on germination, growth, water status, proline content and mineral accumulation of seedlings of *Aegiceras corniculatum* (L.) Blanco (Myrsinaceae). NaCl was added to the soil and salinity was maintained at 0.3, 3.9, 7.9, 12.1, 16.1, 19.7, 24.0, 28.0, 32.0, 36.0 and 40.0 dSm⁻¹. This plant is highly salt tolerant at germination stage. Growth of seedlings was significantly promoted by low salinity and their optimum growth was obtained at 24.0 dSm⁻¹. Moreover, higher salinities inhibited plant growth. Water potential of tissues became significantly more negative with increase in salinity. Proline content in tissues significantly increased as salinity increased. Concentration of Na in tissues significantly increased, whereas K, Ca, N and P content decreased with increase in salinity. Changes in tissues and whole-plant accumulation patterns of other nutrients, as well as possible mechanisms for avoidance of Na toxicity in this plant in response to salinity, are discussed.

Key words: *Aegiceras corniculatum*; Mineral accumulation; Proline content; Propagule germination; Seedling growth; Water status

Introduction

Mangroves inhabit intertidal zones with high salinity (Shan et al. 2008) and are able to tolerate a large range of salinities under natural conditions (Suarez et al. 1998). The growth and physiological mechanisms of mangroves differ in nature due to their complexity of structure and differences in flooding regime, tidal inundation, rapid influx of extra nutrients as well as type of soil (Clough 1984, Naidoo 1987). They possess a variety of adaptations to extreme environmental stresses such as (i) salt exclusion by root ultrafiltration (Scholander 1968), (ii) salt secretion via glands (Roth 1992), (iii) ion accumulation in leaf cells (Popp 1994), (iv) leaf succulence (Roth 1992) and (v) accumulating organic acids as osmotica to counter toxic effects of salinity (Popp 1984). Like other halophytes, mangroves decrease their water and osmotic potentials to maintain turgor at high salinity (Naidoo 1987, Khan et al. 2000a, b). Salinity required for optimal growth varies from 10% to 50% seawater (Downton 1982, Clough 1984, Naidoo 1987, Lin & Sternberg 1992, 1995) and a decline in their optimal growth is obtained with a further increase in salinity. Similarly, lower water potential and accumulation of inorganic ions are the results from extreme saline environments for most of the plants (Ball & Farquhar 1984, Naidoo 1987).

The mangroves are found in sandy and muddy intertidal zones of Arabian Sea along semi-arid region of Saurashtra and contiguous saline desert of Kutch in Gujarat State of India. Overexploitation, pollution and other biotic stresses have drastically reduced and fragmented the mangrove forests. Little information exists on the salt tolerance of mangroves of Saurashtra and Kutch and this information is crucial to the success of restoration effort in the region. *Aegiceras corniculatum* is one of the most common tree species along the coastline in Gujarat. The present study is designed to investigate the salt tolerance of *Aegiceras corniculatum* (L.) Blanco (Myrsinaceae) growing along the coasts of Saurashtra and Kutch in Gujarat.

Material and Methods

Study area

The present study was carried out in a greenhouse

of the botanical garden of Saurashtra University at Rajkot (22°18' N Lat, 70°56' E Long) in Gujarat. For propagule germination and growth of seedlings, the top 15 cm black-cotton soil (Vertisol), which is predominant in Saurashtra region of Gujarat, was collected from an agricultural field. This soil is a clayey loam containing 19.6% sand, 20.3% silt and 60.1% clay. The available soil water between wilting coefficient and field capacity ranged from 18.3% to 35.0%, respectively. The total organic carbon content was 1.3% and pH was 7.2. The electrical conductivity of soil was 0.3 dSm⁻¹. Nitrogen, phosphorus, potassium, calcium and sodium contents were 0.15%, 0.05%, 0.03%, 0.05% and 0.002%, respectively. This soil is fertile and fit for intensive agriculture. Physical and chemical properties of soil are given earlier (Pandya et al. 2004).

Salinisation of soil

Surface soil was collected, air dried and passed through a 2mm mesh screen. Eleven lots of soil, of 100 kg each, were separately spread, about 50 mm thick, over polyethylene sheets. Sodium chloride (NaCl) amounting to 280, 690, 1410, 1900, 2400, 2900, 3300, 3800, 4200 and 4700 g was then thoroughly mixed with soil of ten lots, respectively to give electrical conductivities of 3.9, 7.9, 12.1, 16.1, 19.7, 24.0, 28.0, 32.0, 36.0 and 40.0 dSm⁻¹. There was no addition of NaCl to seventh lot of soil that served as control. Seawater salinity at Jamnagar coast in Saurashtra varies from 45 to 48 dSm⁻¹ during the rainy (monsoon) season. Thus, soil salinity in the present investigation was not maintained above 40 dSm⁻¹. The electrical conductivity of control soil was 0.3 dSm⁻¹ and this value was approximately equal to 3.0 mM salinity. For the measurement of electrical conductivity a soil suspension was prepared in distilled water at 1:2 soil:water ratio. The suspension was shaken and allowed to stand overnight. Thereafter, electrical conductivity of the supernatant solution was determined with a conductivity meter.

Seedling emergence

Twenty polyethylene bags for each level of soil salinity were each filled with 5 kg of soil. Tap water was added to each bag until water level was 2 cm high above the soil surface. Bags were kept in an uncontrolled greenhouse under natural tem-

perature and light. Propagules of *A. corniculatum* were collected from Jamnagar coast. Ten propagules were sown (propagules were gently pushed to one-third of their length into the soil) in each bag on 27th June 2008. Tap water was added daily to compensate evapotranspiration loss. Emergence of seedlings was recorded daily over a period of 30 days. A linear model was fitted to cumulative proportion of propagule germination and increasing soil salinity using the expression:

$$\text{Sin}^{-1} \sqrt{P} = \beta_0 + \beta_1 X$$

where, $\text{Sin}^{-1} \sqrt{P}$ is the proportion of cumulative seed germination, X is soil salinity and β_0 and β_1 are constants. Salt concentration at which seed germination was reduced to 50% (SG_{50}) was estimated using the model.

Seedling growth

For the growth studies, two seedlings that emerged first were left in each of 20 bags at each level of salinity and others were uprooted. Seedlings grown in soils at 0.3, 3.9, 7.9, 12.1, 16.1, 19.7, 24.0, 28.0, 32.0, 36.0 and 40.0 dSm^{-1} salinities exhibited emergence of the second leaf after 9, 10, 10, 12, 12, 12, 12, 12, 12 and 13 days, respectively. Emergence of the second leaf confirmed the establishment of seedlings. Following emergence of the second leaf, one seedling having better vigor was allowed to grow in each bag and another seedling was further uprooted. Thus twenty replicates factorialized with eleven grades of soil (0.3, 3.9, 7.9, 12.1, 16.1, 19.7, 24.0, 28.0, 32.0, 36.0 and 40.0 dSm^{-1}) were prepared. This gave a total of 220 bags, which were arranged in twenty randomized blocks. Seedlings were watered daily to maintain water level above the soil surface and experiment was terminated after six months. The mean maximum temperature of the greenhouse during the course of study decreased from $35.6 \pm 0.57^\circ\text{C}$ in June to $30.8 \pm 0.35^\circ\text{C}$ in August 2008 and further increased to $35.9 \pm 0.69^\circ\text{C}$ in October. Following this period, the mean maximum temperature gradually decreased to $27.8 \pm 0.32^\circ\text{C}$ in January 2009. Seedlings contained in 20 bags at each salinity level were washed with tap water to remove salt on leaves and soil particles adhered to roots. Roots of *A. corniculatum* are adventitious and arise from the root stock. Morphological characteristics of each propagule were recorded. Shoot height and

length of adventitious roots of the propagule were measured. Leaf area was marked out on graph paper. Fresh and dry weights of tissues (leaves, stems and roots) were determined. Water content (gg^{-1} dry weight) in plant tissues was calculated using fresh and dry weight values. Data recorded for morphological characteristics, dry weight and water content of different components were analyzed by one-way ANOVA to assess the effect of salinity on plant growth. In addition, data for growth parameters and water content in tissues were fitted to second order polynomial equation by computer curvilinear method and correlations between examined parameters and soil salinity were computed.

Determination of water potential and proline content

Ten additional plants grown in soil at each level of salinity were used for measurement of water potential and proline determination in plant tissues. Water potential of leaves, stems and roots was measured by Dewpoint Potential Meter WP4 following Patel et al. (2009). All the measurements were taken between 8 to 10.30 A.M. Concentration of proline in plant tissues was estimated following Bates et al. (1973). Extract of 0.5 g fresh plant material with aqueous sulphosalicylic acid was prepared. The extracted proline was made to react with ninhydrin to form chromophore and read at 520 nm. Data were analyzed by one-way ANOVA. Correlations and linear regression equations for water potential and proline content of tissues with salinity were determined.

Mineral analyses of plant materials

Mineral analyses were performed on leaves, stems and root tissues. Plant parts of the seedlings grown in soil at same level of salinity were pooled separately. Plant samples were ground using mortar and pestle. Three subsamples of plant tissues were analyzed. Total nitrogen was determined by Kjeldahl method and phosphorus content estimated by the chlorostannous molybdophosphoric blue colour method in sulphuric acid (Piper 1944). Concentrations of Ca, Mg, Na, K, Zn, Fe, Mn and Cu were determined by Shimadzu double beam atomic absorption spectrophotometer AA-6800 after triacid ($\text{HNO}_3:\text{H}_2\text{SO}_4:\text{HClO}_4$ in the ratio of 10:1:4) digestion. Mineral data were analysed by

one-way ANOVA. Correlations and linear regression equations between mineral content and salt concentrations were determined.

Results

Effect of salinity on propagule germination

Propagules began to germinate 3 days after sowing and 100% propagules germinated over a period of 18 days under control (0.3 dSm⁻¹ salinity) conditions. (Fig. 1). Propagule germination in saline soils was recorded 4-7 days after sowing. Germination lasted for 19, 19, 20, 20, 20, 20, 20, 20 and 20 days in soils with 3.9, 7.9, 12.1, 16.1, 19.7, 24.0, 28.0, 32.0, 36.0 and 40.0 dSm⁻¹ salinities, respectively and corresponding germination was 99.6%, 98.8%, 98.4%, 97.6%, 97.2%, 96.8%, 96.4%, 96.0%, 94.0% and 90.4%. There was a significant reduction in germination of propagules ($p < 0.01$) with increasing salt stress. A negative relationship between proportion of cu-




Figura 1. Emergencia acumulada de semillas de *Aegiceras corniculatum* en respuesta a la salinidad del suelo. 0.3 dS m⁻¹ (●), 3.9 dSm⁻¹ (○), 7.9 dSm⁻¹ (▲), 12.1 dSm⁻¹ (Δ), 16.1 dSm⁻¹ (◆), 19.7 dSm⁻¹ (◇), 24.0 dSm⁻¹ (■), 28.0 dSm⁻¹ (□), 32.0 dSm⁻¹ (•), 36.0 dSm⁻¹ (◊) y 40.0 dSm⁻¹ (◄). Las barras de error representan ES.

Figure 1. Cumulative emergence of seedlings of *Aegiceras corniculatum* in response to soil salinity. 0.3 dS m⁻¹ (●), 3.9 dSm⁻¹ (○), 7.9 dSm⁻¹ (▲), 12.1 dSm⁻¹ (Δ), 16.1 dSm⁻¹ (◆), 19.7 dSm⁻¹ (◇), 24.0 dSm⁻¹ (■), 28.0 dSm⁻¹ (□), 32.0 dSm⁻¹ (•), 36.0 dSm⁻¹ (◊) and 40.0 dSm⁻¹ (◄). Error bars represent SE.

mulative propagule germination and concentration of salt was obtained according to the following expression:

$$Y = 95.700 - 0.500X \quad (R^2_{Adj} = 0.969, p < 0.01)$$

where, Y is arcsine (degrees) of proportion of cumulative seed germination and X is salt concentration.

Effect of salinity on stem and root elongation and leaf expansion

Salinity significantly stimulated ($p < 0.01$) stem and root elongation until 24.0 dSm⁻¹, but extension growth declined with further increases in salinity (Table 1). In general, stem height was manifold greater than root length under the control and saline conditions. There was a quadratic relationship for shoot height and root length with salt concentration ($p < 0.01$). Leaf area significantly increased ($p < 0.01$) until 24.0 dSm⁻¹ salinity, but decreased at higher salinities (Table 1). A quadratic relationship was obtained between leaf area and salt concentration in soil ($p < 0.01$).

Effect of salinity on dry weight

Dry weight of leaves, stems, shoots (leaves + stems) and roots was significantly promoted ($p < 0.01$) by salinity until 24.0 dSm⁻¹, but it declined with further increases in salinity (Table 1). There was a quadratic relationship between salt concentration in soil and dry weight of tissues ($p < 0.01$). Root/shoot dry weight ratio was 0.10 for plants grown in control soil and it did not change with increasing soil salinity.

Effect of salinity on water content and water potential of tissues

Water content (gg⁻¹ dry weight) significantly increased ($p < 0.01$) in leaves, stems and roots until 24.0 dSm⁻¹ salinity, but declined with further increases in salinity (Fig. 2A). A quadratic relationship was obtained between soil salinity and water content of leaves ($r = 0.465$, $p < 0.01$), stems ($r = 0.565$, $p < 0.01$) and roots ($r = 0.382$, $p < 0.01$). Tissues significantly differed ($p < 0.01$) in their water content. Moreover, maximum water content was in leaves and roots, and minimum in stems. Tissues according to their water content can be arranged into the following decreasing order: leaves = roots > stems. Water potential significantly became more negative in tissues ($p < 0.01$) as soil sa-

Salinity (dSm ⁻¹)	Stem height (cm)	Root length (cm)	Leaf area (cm ² plant ⁻¹)	Leaf weight (mg plant ⁻¹)	Stem weight (mg plant ⁻¹)	Shoot weight (leaf+stem) (mg plant ⁻¹)	Root weight (mg plant ⁻¹)	Root/Shoot dry weight ratio
0.3	21.0 ± 0.6	4.5 ± 0.2	25.7 ± 2.0	424.5 ± 22.4	1826.5 ± 94.8	2251.0 ± 102.3	232.5 ± 14.3	0.1 ± 0.0
3.9	22.9 ± 0.5	5.6 ± 0.2	27.9 ± 1.2	503.0 ± 24.8	2101.0 ± 74.5	2604.0 ± 84.2	271.0 ± 16.1	0.1 ± 0.0
7.9	23.1 ± 0.5	6.0 ± 0.1	28.9 ± 1.1	529.5 ± 27.2	2151.0 ± 80.6	2680.5 ± 97.1	289.0 ± 16.1	0.1 ± 0.0
12.1	24.1 ± 0.5	6.5 ± 0.2	35.5 ± 1.3	617.5 ± 33.3	2328.5 ± 101.8	2946.0 ± 119.8	299.5 ± 19.6	0.1 ± 0.0
16.1	25.5 ± 0.4	7.8 ± 0.2	41.1 ± 1.4	665.5 ± 25.9	2486.5 ± 85.7	3152.0 ± 98.4	336.0 ± 21.6	0.1 ± 0.0
19.7	27.4 ± 0.4	11.0 ± 0.4	47.8 ± 1.4	795.0 ± 27.6	2645.5 ± 82.6	3440.5 ± 97.9	361.5 ± 15.8	0.1 ± 0.0
24.0	29.0 ± 0.2	11.8 ± 0.2	54.1 ± 1.2	1002.0 ± 37.4	2779.5 ± 58.9	3781.5 ± 87.3	394.5 ± 15.4	0.1 ± 0.0
28.0	23.9 ± 0.3	10.1 ± 0.3	45.5 ± 1.2	771.5 ± 22.9	2558.5 ± 71.2	3330.0 ± 85.2	347.0 ± 16.2	0.1 ± 0.0
32.0	23.1 ± 0.3	6.1 ± 0.2	39.4 ± 1.2	736.5 ± 36.9	2341.0 ± 60.4	3077.5 ± 82.3	324.0 ± 12.5	0.1 ± 0.0
36.0	21.9 ± 0.4	5.8 ± 0.2	37.4 ± 0.6	675.5 ± 37.7	2175.5 ± 70.9	2851.0 ± 87.2	294.0 ± 19.7	0.1 ± 0.0
40.0	21.1 ± 0.4	5.1 ± 0.2	31.8 ± 1.1	621.5 ± 27.7	1939.0 ± 72.2	2560.5 ± 87.5	266.5 ± 11.3	0.1 ± 0.0
α	20.350	3.310	20.260	356.900	1746.000	2103.000	218.500	NS
β	-0.014	-0.013	-0.047	-0.726	-1.883	-2.610	-0.281	NS
γ	0.588	0.579	2.189	35.840	80.490	116.300	12.450	NS
r	0.671	0.747	0.742	0.659	0.607	0.666	0.494	NS
LSD _{0.05}	1.4	0.8	4.2	97.8	256.7	307.5	53.9	NS

Relationship is significant at $p < 0.01$; NS = Non significant.

Tabla 1. Efecto de la salinidad del suelo en las características de la hoja, tallo y raíz de *Aegiceras corniculatum* indicado por la media \pm SEM y las constantes de la ecuación de regresión.

Table 1. Effect of soil salinity on leaf, stem, shoot and root characteristics of *Aegiceras corniculatum* as indicated by mean \pm SEM and regression equation constants.

linity increased (Fig. 2B). There was a negative relationship between soil salinity and water potential of leaves ($r = -0.953$, $p < 0.01$), stems ($r = -0.926$, $p < 0.01$), and roots ($r = -0.912$, $p < 0.01$). Tissues significantly differed ($p < 0.01$) in their water potential. Tissues according to their negative water potential values (low to high negative values) can be arranged into the following decreasing order: roots > stems > leaves.

Effect of salinity on proline content of tissues

Proline content ($\mu\text{mol/g}$ FW material) significantly increased ($p < 0.01$) in leaves, stems, and root tissues with increase in soil salinity (Fig. 2C). Tissues according to their proline content can be arranged into the following decreasing order: stems > leaves > roots. There was a positive relationship between salt concentration and proline content of leaves ($r = 0.953$, $p < 0.01$), stems ($r = 0.952$, $p < 0.01$) and roots ($r = 0.959$, $p < 0.01$). A

significant inverse relationship was obtained between water potential (more negative values) and proline content of leaves ($r = -0.911$, $p < 0.01$), stems ($r = -0.851$, $p < 0.01$) and roots ($r = -0.852$, $p < 0.01$).

Effect of salinity on mineral accumulation

Sodium content significantly increased ($p < 0.01$) in leaves, stems and roots with increase in salinity (Table 2). A positive relationship was obtained between Na content of tissues and salt stress ($p < 0.01$). K content significantly decreased ($p < 0.01$) in leaves, stems and root tissues, in response to increase in salinity. There was a negative relationship between K content of tissues and salt concentration in soil ($p < 0.01$). The Na/K ratio significantly increased ($p < 0.01$) in leaves, stems and root tissues in response to increasing soil salinity. There was positive relationship ($p < 0.01$) between Na/K ratio in tissues and salt stress.

Concentration of N, P and Ca significantly de-




Figura 2. Efecto de la salinidad del suelo en: **A.** contenido de agua; **B.** potencial hídrico; **C.** contenido de prolina en hojas (●), tallos (○) y raíces (▲) de *Aegiceras corniculatum*. Las barras de error representan SE.

Figure 2. Effect of salinisation of soil on: **A.** water content; **B.** water potential; **C.** proline content of leaves (●), stems (○) and roots (▲) of *Aegiceras corniculatum*. Error bars represent SE.

creased ($p < 0.01$) in leaves, stems and roots in response to increasing salt concentration in soil. A negative relationship was obtained between N, P and Ca content in tissues and salt concentration ($p < 0.01$). Mg content in tissues did not change with increase in salt concentration in soil. The concentration of Zn, Mn and Fe significantly increased ($p < 0.01$) in leaves, stems and roots in response to increase in salt-stress (Table 2). A positive relationship was obtained between soil salinity and Zn, Mn and Fe content in tissues ($p < 0.01$). There was a significant decrease ($p < 0.01$) in the concentration of Cu in leaves, stems and roots in response to increase in salinity. A negative relationship was obtained between salt concentration and Cu content in tissues ($p < 0.01$).

Discussion

Optimum germination (100%) was obtained in non-saline control soil and increasing salinity delayed and reduced germination of *A. corniculatum*. Similar results have been reported by others for seed germination of halophytes (Khan & Weber 1986, Ungar 1996, Katembe et al. 1998, Gulzar & Khan 2001, Khan 2002, Li et al. 2002). Salt can affect seed germination either by osmotic effect, restricting the supply of water to embryo, (Agboola 1998, Pujol et al. 2000) or by ionic effect, causing specific injury through ions to the metabolic machinery, (Pollack & Waisel 1972, Mohammad & Sen 1990). Propagule germination of *A. corniculatum* could be reduced to 50% (SG_{50}) at salinity of 91.4 dSm⁻¹, whereas SG_{50} for *Cassia montana*, a halophyte tree in coastal region of Saurashtra, was obtained at 6.0 dSm⁻¹ (Patel & Pandey 2007). That would suggest that this plant species is highly salt tolerant at germination stage. *A. corniculatum* is characterized by viviparous germination. Adaptation of viviparous propagules to saline environments actually starts when they are still attached to the mother tree by continuously absorbing salt from the tree or by a desalinating process (Joshi et al. 1972, Zheng et al. 1999). Under natural conditions, germination of propagules of *A. corniculatum* occurs during the season with high precipitation when salinity level of seawater is usually reduced. It appears that *A. corniculatum* invades coastline along the Arabian Sea because of its high salt tolerance at seed germination stage.

Growth of *A. corniculatum* seedlings was stimulated by low salinity and their optimum growth was at 24.0 dSm⁻¹. Similar results have been reported for halophytes that have optimal growth in the presence of salt (Naidoo & Raghunanan 1990, Ayala & O'Leary 1995, Khan et al. 2000a, Patel & Pandey 2007). Soil salinity at 24.0 dSm⁻¹ was equal to 50% seawater treatment because salinity of seawater during the rainy season at Jamnagar coast is about 48.0 dSm⁻¹. Result does not agree with some other studies in which mangrove species showed their best growth at 25% seawater concentration (Downton 1982, Clough 1984, Naidoo 1987). However, Karim & Karim (1993) reported that the growth for *Avicennia marina* from Bangladesh coast was obtained at 50% seawater. High media salinity affects plant

Tissue	Salinity (dSm ⁻¹)	N (mg g ⁻¹)	P (mg g ⁻¹)	K (mg g ⁻¹)	Na (mg g ⁻¹)	Ca (mg g ⁻¹)	Mg (mg g ⁻¹)	Na/K ratio	Zn (µg g ⁻¹)	Cu (µg g ⁻¹)	Mn (µg g ⁻¹)	Fe (µg g ⁻¹)
Leaf	0.3	24.0 ± 1.4	2.2 ± 0.3	11.4 ± 0.6	7.1 ± 0.8	12.6 ± 0.2	1.4 ± 0.0	0.6 ± 0.1	29.6 ± 2.8	16.3 ± 0.7	85.7 ± 5.2	267.0 ± 27.2
	3.9	24.0 ± 1.5	2.2 ± 0.1	11.0 ± 0.8	7.6 ± 1.0	12.2 ± 0.7	1.4 ± 0.1	0.7 ± 0.1	33.1 ± 2.6	15.4 ± 1.1	91.3 ± 5.4	331.2 ± 16.9
	7.9	22.0 ± 1.7	2.2 ± 0.1	10.3 ± 0.9	9.2 ± 1.5	11.7 ± 0.7	1.3 ± 0.1	0.9 ± 0.1	38.4 ± 2.8	14.4 ± 0.6	97.0 ± 8.4	347.3 ± 15.8
	12.1	21.0 ± 1.3	2.1 ± 0.1	10.0 ± 0.9	10.7 ± 0.6	11.1 ± 0.5	1.3 ± 0.1	1.1 ± 0.1	39.3 ± 2.0	14.0 ± 0.6	103.2 ± 7.3	365.7 ± 19.1
	16.1	20.0 ± 1.3	2.0 ± 0.2	9.8 ± 0.4	10.9 ± 1.1	10.8 ± 0.4	1.3 ± 0.1	1.1 ± 0.1	42.2 ± 1.6	13.0 ± 0.5	109.2 ± 7.1	392.3 ± 20.7
	19.7	19.0 ± 1.5	1.9 ± 0.3	9.0 ± 0.7	11.4 ± 1.0	10.2 ± 0.6	1.2 ± 0.1	1.3 ± 0.1	46.3 ± 2.4	13.0 ± 0.6	114.1 ± 6.8	441.1 ± 14.6
	24.0	19.0 ± 1.0	1.8 ± 0.1	8.5 ± 1.0	11.7 ± 0.8	9.9 ± 0.9	1.2 ± 0.0	1.4 ± 0.3	49.1 ± 2.1	12.3 ± 0.9	116.4 ± 6.5	422.9 ± 13.2
	28.0	19.0 ± 1.2	1.7 ± 0.2	8.3 ± 0.8	11.7 ± 0.8	9.9 ± 0.6	1.2 ± 0.1	1.4 ± 0.2	51.2 ± 2.7	11.4 ± 0.7	122.3 ± 5.8	460.3 ± 18.8
	32.0	18.0 ± 1.0	1.5 ± 0.2	7.8 ± 1.1	12.2 ± 0.9	9.3 ± 0.9	1.2 ± 0.1	1.6 ± 0.3	55.4 ± 2.2	11.3 ± 1.3	123.0 ± 7.1	523.1 ± 19.3
	36.0	17.0 ± 1.0	1.3 ± 0.1	7.2 ± 0.8	12.6 ± 0.7	8.9 ± 0.6	1.1 ± 0.0	1.8 ± 0.3	57.1 ± 2.5	10.4 ± 1.2	127.0 ± 7.6	587.7 ± 21.8
40.0	17.0 ± 1.3	1.2 ± 0.2	7.1 ± 0.9	12.7 ± 1.0	8.7 ± 1.0	1.1 ± 0.1	1.9 ± 0.4	59.3 ± 2.8	10.0 ± 1.4	129.7 ± 5.9	683.3 ± 14.8	
α	23.656	2.384	11.360	7.994	12.454	NS	NS	0.624	30.725	15.916	88.677	259.941
β	-0.183	-0.027	-0.112	0.135	-0.098	NS	NS	0.032	0.742	-0.152	1.107	8.784
r	-0.764	-0.788	-0.769	0.747	-0.783	NS	NS	0.813	0.934	-0.822	0.819	0.933
LSD_{0.05}	3.970	0.550	2.490	2.930	2.070	NS	NS	0.610	7.430	2.800	20.370	56.940
Stem	0.3	22.0 ± 1.0	2.0 ± 0.1	9.1 ± 0.5	10.5 ± 0.5	11.9 ± 0.5	1.3 ± 0.0	1.2 ± 0.1	35.4 ± 4.3	13.3 ± 0.9	59.0 ± 4.9	410.0 ± 20.5
	3.9	21.0 ± 0.7	1.9 ± 0.2	9.0 ± 0.5	10.9 ± 0.3	11.8 ± 0.6	1.3 ± 0.0	1.2 ± 0.1	37.1 ± 3.3	12.7 ± 0.5	65.1 ± 5.8	460.1 ± 21.1
	7.9	21.0 ± 1.0	1.9 ± 0.2	8.7 ± 0.2	11.2 ± 0.7	11.0 ± 0.7	1.3 ± 0.0	1.3 ± 0.1	43.1 ± 2.2	12.4 ± 0.5	69.2 ± 4.5	513.1 ± 15.5
	12.1	20.0 ± 1.2	1.8 ± 0.1	8.3 ± 1.0	11.3 ± 0.4	10.9 ± 0.7	1.2 ± 0.1	1.4 ± 0.2	48.3 ± 2.3	11.3 ± 0.6	77.0 ± 4.7	563.7 ± 17.7
	16.1	19.0 ± 1.0	1.8 ± 0.1	7.9 ± 0.3	11.7 ± 0.4	10.5 ± 0.6	1.2 ± 0.1	1.5 ± 0.1	50.2 ± 2.7	11.0 ± 0.3	79.6 ± 6.5	661.5 ± 16.4
	19.7	18.0 ± 1.1	1.7 ± 0.2	7.7 ± 0.5	11.9 ± 0.2	9.7 ± 0.7	1.2 ± 0.1	1.6 ± 0.1	53.1 ± 2.3	10.8 ± 0.6	83.7 ± 6.5	747.7 ± 16.2
	24.0	18.0 ± 1.0	1.6 ± 0.1	7.3 ± 0.3	12.2 ± 0.2	9.5 ± 0.9	1.1 ± 0.0	1.7 ± 0.1	55.1 ± 3.8	10.3 ± 0.5	95.6 ± 4.9	763.3 ± 19.4
	28.0	18.0 ± 1.1	1.5 ± 0.3	6.8 ± 0.6	12.6 ± 0.3	9.5 ± 0.5	1.1 ± 0.1	1.9 ± 0.1	57.2 ± 3.0	10.0 ± 0.4	102.6 ± 7.1	787.1 ± 20.1
	32.0	17.0 ± 1.0	1.4 ± 0.1	6.6 ± 0.5	12.7 ± 0.5	8.9 ± 0.6	1.1 ± 0.1	1.9 ± 0.2	59.4 ± 2.0	10.0 ± 0.7	106.3 ± 4.3	801.0 ± 15.3
	36.0	16.0 ± 0.6	1.3 ± 0.2	6.4 ± 0.4	12.9 ± 0.5	8.4 ± 0.9	1.1 ± 0.1	2.0 ± 0.1	64.6 ± 1.9	9.8 ± 0.6	111.5 ± 10.2	821.3 ± 17.0
40.0	16.0 ± 1.0	1.2 ± 0.1	6.4 ± 0.7	13.1 ± 0.7	8.3 ± 0.6	1.0 ± 0.1	2.1 ± 0.2	66.3 ± 1.5	9.3 ± 0.6	116.3 ± 5.7	866.0 ± 15.4	
α	21.785	2.073	9.166	10.615	11.938	NS	NS	1.124	36.471	12.917	58.117	438.467
β	-0.152	-0.021	-0.076	0.064	-0.094	NS	NS	0.024	0.767	-0.096	1.484	11.690
r	-0.802	-0.761	-0.770	0.781	-0.770	NS	NS	0.852	0.918	-0.813	0.904	0.957
LSD_{0.05}	2.970	0.470	1.680	1.380	2.080	NS	NS	0.400	8.440	1.750	18.610	54.060

Relationship is significant at p < 0.01. NS = non significant.

Tabla 2. Efecto de la salinidad del suelo en el contenido de nutrientes de los tejidos (hoja, tallo y raíz) de *Aegiceras corniculatum* indicado como media ± SEM and constantes de la ecuación de regresión.

Table 2. Effect of soil salinity on nutrient content of tissues (leaf, stem and root) of *Aegiceras corniculatum* as indicated by mean ± SEM and regression equation constants.

Tissue	Salinity (dSm ⁻¹)	N (mg g ⁻¹)	P (mg g ⁻¹)	K (mg g ⁻¹)	Na (mg g ⁻¹)	Ca (mg g ⁻¹)	Mg (mg g ⁻¹)	Na/K ratio	Zn (µg g ⁻¹)	Cu (µg g ⁻¹)	Mn (µg g ⁻¹)	Fe (µg g ⁻¹)
	0.3	21.0 ± 1.2	1.4 ± 0.1	6.8 ± 0.2	10.6 ± 0.6	10.7 ± 0.7	1.2 ± 0.1	1.6 ± 0.1	50.3 ± 2.1	11.4 ± 0.2	41.9 ± 2.9	991.7 ± 29.4
	3.9	20.0 ± 0.6	1.3 ± 0.1	6.5 ± 0.3	10.9 ± 0.6	10.6 ± 0.3	1.2 ± 0.1	1.7 ± 0.1	51.7 ± 3.5	11.1 ± 0.1	43.1 ± 3.0	1012.3 ± 20.3
	7.9	20.0 ± 1.2	1.3 ± 0.1	6.5 ± 0.2	11.2 ± 0.5	10.4 ± 0.2	1.2 ± 0.1	1.7 ± 0.1	54.9 ± 2.9	10.8 ± 0.1	46.5 ± 3.0	1101.3 ± 22.7
	12.1	19.0 ± 1.0	1.2 ± 0.2	6.1 ± 0.1	11.3 ± 0.4	10.1 ± 0.5	1.1 ± 0.1	1.8 ± 0.1	57.3 ± 2.5	10.7 ± 0.2	48.7 ± 3.7	1237.6 ± 21.5
	16.1	18.0 ± 1.4	1.0 ± 0.1	6.0 ± 0.3	11.7 ± 1.0	9.9 ± 0.6	1.1 ± 0.1	2.0 ± 0.1	61.7 ± 3.0	10.5 ± 0.2	56.4 ± 2.6	1504.8 ± 16.9
	19.7	17.0 ± 1.0	1.0 ± 0.1	5.8 ± 0.2	11.7 ± 0.5	9.1 ± 0.5	1.1 ± 0.0	2.0 ± 0.1	63.4 ± 2.6	10.5 ± 0.2	59.9 ± 3.8	1696.5 ± 25.8
	24.0	17.0 ± 0.8	0.9 ± 0.1	5.8 ± 0.1	12.1 ± 0.4	9.1 ± 0.7	1.1 ± 0.1	2.1 ± 0.1	67.8 ± 2.8	10.3 ± 0.2	64.5 ± 2.8	1882.0 ± 20.7
Root	28.0	16.0 ± 1.1	0.9 ± 0.0	5.7 ± 0.2	12.5 ± 0.7	8.7 ± 0.4	1.0 ± 0.0	2.2 ± 0.2	69.1 ± 2.1	10.2 ± 0.2	66.1 ± 2.3	2046.3 ± 27.1
	32.0	16.0 ± 0.7	0.8 ± 0.1	5.6 ± 0.3	13.1 ± 0.4	8.6 ± 0.7	1.0 ± 0.1	2.3 ± 0.1	72.7 ± 2.0	10.1 ± 0.3	68.2 ± 3.2	2156.5 ± 28.7
	36.0	15.0 ± 1.0	0.8 ± 0.1	5.5 ± 0.3	13.6 ± 0.3	8.1 ± 0.3	1.0 ± 0.1	2.5 ± 0.2	75.7 ± 2.6	10.0 ± 0.4	69.1 ± 2.9	2380.0 ± 23.1
	40.0	15.0 ± 1.0	0.8 ± 0.1	5.5 ± 0.2	13.8 ± 0.6	7.9 ± 0.6	1.0 ± 0.0	2.5 ± 0.1	76.1 ± 1.9	9.8 ± 0.3	72.1 ± 2.8	2454.3 ± 32.8
	α	20.787	1.339	6.634	10.440	10.870	NS	1.551	49.656	11.195	41.340	860.545
	β	-0.158	-0.015	-0.032	0.080	-0.075	NS	0.024	0.702	-0.036	0.826	40.897
	r	-0.802	-0.775	-0.751	0.770	0.774	NS	0.857	0.921	-0.776	0.911	0.990
	LSD_{0.05}	3.100	0.330	0.730	1.760	1.620	NS	0.390	7.850	0.760	9.190	75.450

Relationship is significant at $p < 0.01$, NS = non significant.

Tabla 2 (continúa). Efecto de la salinidad del suelo en el contenido de nutrientes de los tejidos (hoja, tallo y raíz) de *Aegiceras corniculatum* indicado como media ± SEM and constantes de la ecuación de regresión.

Table 2 (continues). Effect of soil salinity on nutrient content of tissues (leaf, stem and root) of *Aegiceras corniculatum* as indicated by mean ± SEM and regression equation constants.

growth due to low water potential, ion toxicities, nutrient deficiencies or a combination of all these factors (Khan et al. 2000 a). Patel and Pandey (2007) reported that seedlings of *Cassia montana*, a halophyte tree in the coastal area of Saurashtra, exhibited optimum growth at 7.9 dSm⁻¹ salinity. Evidently, *A. corniculatum* can be grouped among highly salt tolerant plants. Root/shoot dry weight ratio did not change with increases in salinity suggesting that there was resemblance in shoot and root growth of *A. corniculatum*.

Water content of tissues increased until 24.0 dSm⁻¹ and then declined with increased salinity. Similar result has been reported for *Suaeda fruticosa* (Khan et al. 2000a). Halophytes are characterized by their capacity to adjust tissue water potential to a level that is more negative than that of the soil water potential of the habitat in which they are growing (Ungar 1991). Mangroves lower tissue osmotic potential through the net accumulation of solutes in response to salinity and water deficits (Turner & Jones 1993, Suarez et al. 1998). *A. corniculatum* showed a progressive decrease in water potential of leaves and stems with an increase in salinity, indicating that it follows an osmoconformer strategy to maintain its osmotic balance.

The salt secreting *A. corniculatum* had high concentration of Na in tissues when grown in soils with increasing salinity. There are evidences that halophytes and glycophytes accumulate NaCl in vacuoles (Flowers et al. 1977). High internal salt concentrations provide potential benefits to mangrove plants growing under conditions where soil osmotic potential is more negative than that of seawater because of high soil salinity (Ungar 1991). Increased salt content lowers the internal water potential which is required to permit water uptake. Na⁺ and Cl⁻ accumulated in leaf tissues provide osmotic adjustment and turgor to maintain growth (Yeo 1983). The cation K⁺ is essential for cell expansion, osmo-regulation and cellular and whole-plant homeostasis (Schachtman et al. 1997). High stomatal K⁺ requirement is reported for photosynthesis (Chow et al. 1990). The role of K⁺ in response to salt stress is also well documented, where Na⁺ depresses K⁺ uptake (Fox & Guerinot 1998). In the present study, significant decrease of K⁺ content in all the tissues of seedlings with increasing soil salinity suggests that Na⁺ inhibited K⁺ uptake. The Na/K ratio increased

in leaves and stems with increase in salinity suggesting an increase in transportation of Na⁺ from root to shoot. Tattini et al. (1995) reported that Na/K ratio increases in salt tolerant species with increasing salinity in the external medium because mass transport of sodium takes place from root to shoot via the transpiration stream.

The accumulation of compatible solutes is a common response to salinity in higher plants (Stewart & Lee 1974, Storey et al. 1977). These compounds are not toxic to cytoplasmic enzymes functions at high concentrations (Storey et al. 1977), thus help in osmotic adjustment. The increase of proline content in tissues with increase in Na content indicates that higher proline accumulation may contribute to the alleviation of NaCl stress in *A. corniculatum*. Shan et al. (2008) reported that proline is the specific osmolyte for this species. The proline accumulation was greater in stems and leaves (shoot tissues) than that in roots. Munns (2002) concluded that organic solutes are often lower in roots than shoots.

In general, salinity reduces N accumulation in plants (Feigin 1985). This is due to the fact that an increase in chloride uptake and accumulation is mostly accompanied by a decrease in shoot nitrate concentration (Torres & Bingham 1973). The interaction between salinity and P is very complex and there is no clear cut mechanistic explanation for decreased, increased or unchanged P uptake in response to salinisation in different species (Champognol 1979, Grattan & Grieve 1992). However, it is known that P concentration is related to the rate of photosynthesis, since it decreases conversion of fixed carbon into starch (Overlach et al. 1993) and therefore decrease of P in leaves will reduce shoot growth. Calcium is important during salt stress, e.g., in preserving membrane integrity (Rengel 1992), signaling in osmoregulation (Mansfield et al. 1990) and influencing K⁺/Na⁺ selectivity (Cramer et al. 1987). In the present study, there was a significant decrease of Ca²⁺ content in all the tissues with salinisation of soil. As a result, Na⁺ induced Ca²⁺ deficiency in tissues. It is reported that uptake of Ca²⁺ from the soil solution may decrease because of ion interactions, precipitation and increase in ionic strength that reduce the activity of Ca²⁺ (Janzen & Chang 1987). Besides the role of Mg²⁺ in chlorophyll structure and as an enzyme cofactor, another important role of Mg²⁺ in plants is in the export of

photosynthates (Marschner & Cakmak 1989). Results suggested that in extreme saline habitats N, P, K and Ca were limiting factors for the growth of *A. corniculatum* whereas, Mg did not exhibit a similar trend.

It is difficult to suggest mechanistic explanations of salinity influence on micro-elements concentration due to relatively smaller differences between control and salinised tissues (Tozlu et al. 2000). In the present study, accumulation of Zn, Mn and Fe increased, while it decreased for Cu at the whole plant level. Besides, cofactors for enzymes, Fe and Cu are essential for biological redox system, Mn for photolysis of water in photosynthesis and Zn for DNA replication, regulation of gene expression and integrity of biomembranes (Marschner 1995). In addition, high concentration of iron is required for structural and functional integrity of the thylakoid membranes and synthesis of ferredoxin and chlorophyll (Marschner 1995). Decrease of Cu at the whole plant level might limit the growth of plants. Salinity generates an increase in reactive oxygen species (ROS) which have deleterious effects on all metabolisms (Borsani et al. 2001). Superoxide dismutases (SODs) detoxify ROS and may contain Zn, Cu, Mn or Fe as metal component (Slater et al. 2003). Increase of Zn, Mn and Fe content at the whole-plant level might be the requirement of this plant for survival in extreme saline habitats.

Acknowledgement

This study was supported with funds from Departmental Special Assistance provided by the University Grants Commission, New Delhi, Govt. of India.

References

- Agboola DA. 1998. Effect of saline solutions and salt stress on seed germination of some tropical forest tree species. *Tropical Biology* 45: 324-331.
- Ayala F & O'Leary JW. 1995. Growth and physiology of *Salicornia bigelovii* Torr. at suboptimal salinity. *International Journal of Plant Science* 156: 197-205.
- Ball MC & Farquhar JD. 1984. Photosynthetic and stomatal responses of two mangrove species, *Aegiceras corniculatum* and *Avicennia marina* to long term salinity and humidity conditions. *Plant Physiology* 74: 1-6.
- Bates LS, Waldren RP & Teare FD. 1973. Rapid determination of free proline from water stress studies. *Plant and Soil* 39: 205-207.
- Borsani O, Valpuesta V & Botella MA. 2001. Evidence for a role of salicylic acid in the oxidative damage generated by NaCl and osmotic stress in Arabidopsis seedlings. *Plant Physiology* 126: 1024-1030.
- Champagnol F. 1979. Relationship between phosphate nutrition of plants and salt toxicity. *Phosphorus Agriculture* 76: 35-43.
- Chow WS, Ball MC & Anderson JM. 1990. Growth and photosynthetic responses of spinach to salinity, implication of K nutrition for salt tolerance. *Australian Journal of Plant Physiology* 17: 563-578.
- Clough BF. 1984. Growth and salt balance of the mangroves *Avicennia marina* (Forssk.) Vierh. and *Rhizophora stylosa* Griff. in relation to salinity. *Journal of Plant Physiology* 11: 419-430.
- Cramer GR, Lynch J, Lauchli A & Epstein E. 1987. Influx of Na⁺, K⁺ and Ca²⁺ into roots of salt-stressed cotton seedlings. Effects of supplemental Ca. *Plant Physiology* 83: 510-516.
- Downton WJS. 1982. Growth and osmotic relations of the mangrove *Avicennia marina*, as influenced by salinity. *Australian Journal of Plant Physiology* 9: 519-528.
- Feigin A. 1985. Fertilization management of crops irrigated with saline water. *Plant and Soil* 89: 285-299.
- Flowers TJ, Troke PF & Yeo AR. 1977. The mechanism of salt tolerance in halophytes. *Annual Review of Plant Physiology* 28: 89-121.
- Fox TC & Guerinot ML. 1998. Molecular biology of cation transport in plants. *Annual Review of Plant Physiology and Plant Molecular Biology* 49: 669-696.
- Grattan SR & Grieve CM. 1992. Mineral element acquisition and growth response of plants grown in saline environments. *Agriculture Ecosystem and Environment* 38: 275-300.
- Gulzar S & Khan MA. 2001. Seed germination of a halophytic grass *Aeluropus lagopoides*. *Annals of Botany* 87: 319-324.
- Janzen HH & Chang C. 1987. Cation nutrition of barley as influenced by soil solution composition in a saline soil. *Canadian Journal of Soil Science* 67: 619-629.
- Joshi GV, Pimplaskar M & Bhosale LJ. 1972. Physiological studies in germination of mangroves. *Botanica Marina* 45: 91-95.
- Karim J & Karim A. 1993. Effect of salinity on the growth of some mangrove plants in Bangladesh. In: Lieth H & Al Masoom A. (Eds.), *Towards the rational use of high salinity tolerant plants*. Kluwer Academic Publishers, The Netherlands, pp 187-192.
- Katembe J, Ungar IA & Mitchell P. 1998. Effect of salinity on germination and seedlings growth of two *Atriplex* species (Chenopodiaceae). *Annals of Botany* 82: 167-175.
- Khan MA. 2002. Halophyte seed germination: Success and Pitfalls. In: Hegazi AM, El-Shaer HM, El-Demerdashe S, Guirgis RA, Abdel Salam Metwally A, Hasan FA & Khashaba HE. (Eds.), *International symposium on optimum resource utilization in salt affected ecosystems in arid and semi arid regions*.

- Desert Research Centre, Cairo, Egypt, pp 43-50.
- Khan MA & Weber DJ. 1986. Factors influencing seed germination in *Salicornia pacifica* var. *utahensis*. *American Journal of Botany* 73: 1163-1167.
- Khan MA, Ungar IA & Showalter AM. 2000a. The effect of salinity on the growth, water status, and ion content of a leaf succulent perennial halophyte, *Suaeda fruticosa* (L.). *Forssk. Journal of Arid Environments* 45: 73-84.
- Khan MA, Ungar IA & Showalter AM. 2000b. Effects of salinity on growth, water relations and ion accumulation of the subtropical perennial halophyte, *Atriplex griffithii* Moq. var. *stocksii* Boiss. *Annals of Botany* 85: 225-232.
- Li H, Kefu Z & Xiufeng W. 2002. The inhibition of salinity on the germination of halophyte seeds. *Journal of Shandong Agricultural University (natural science)*. 33: 170-173.
- Lin G & Sternberg L. 1992. Effects of growth form, salinity, nutrient and sulfide on photosynthesis, carbon isotope discrimination and growth of red mangrove (*Rhizophora mangle* L.). *Australian Journal of Plant Physiology* 19: 509-517.
- Lin G & Sternberg L. 1995. Variation in propagule mass and its effect on carbon assimilation and seedling growth of red mangrove (*Rhizophora mangle* L.). in Florida, USA. *Journal of Tropical Ecology* 11: 109-119.
- Mansfield TA, Hetherington AM & Atkinson CJ. 1990. Some aspects of stomatal physiology. *Annual Review of Plant Physiology and Plant Molecular Biology* 41: 55-75.
- Marschner H. 1995. *Mineral Nutrition of Higher Plants*. London: Academic Press, pp 889.
- Marschner H & Cakmak I. 1989. High light intensity enhances chlorosis and necrosis in leaves of zinc, potassium and magnesium deficient bean *Phaseolus vulgaris* plants. *Journal of Plant Physiology* 134: 308-315.
- Mohammad S & Sen DN. 1990. Germination behavior of some halophytes in Indian desert. *International Journal of Experimental Biology* 28: 545-549.
- Munns R. 2002. Comparative physiology of salt and water stress. *Plant Cell and Environment* 25: 239-250.
- Naidoo GR & Raghunanan R. 1990. Salt tolerance in the succulent halophyte, *Sarcocornia natalensis*. *Journal of Experimental Botany* 41: 497-502.
- Naidoo G. 1987. Effects of salinity and nitrogen on growth and plant water relations in the mangrove *Avicennia marina* (Forssk.) Vierh. *New Phytologist* 107: 317-326.
- Overlach S, Diekmann W & Raschke K. 1993. Phosphate translocator of isolated guard-cell chloroplasts from *Pisum sativum* L. transport glucose-6-phosphate. *Plant Physiology* 101: 1201-1207.
- Pandya DH, Mer RK, Prajith PK & Pandey AN. 2004. Effect of salt stress and manganese supply on growth of barley seedlings. *Journal of Plant Nutrition* 27: 1361-1379.
- Patel AD & Pandey AN. 2007. Effect of soil salinity on growth, water status and nutrient accumulation in seedlings of *Cassia montana* (Fabaceae). *Journal of Arid Environments* 70: 174-182.
- Patel AD, Jadeja HR & Pandey AN. 2009. Effect of salinisation of soil on growth, water status and nutrient accumulation in seedlings of *Acacia auriculiformis* (Fabaceae). *Journal of Plant Nutrition* (In Press).
- Piper CS. 1944. *Soil and Plant Analysis*. Interscience, New York, pp 368.
- Pollack G & Waisel Y. 1972. Salt secretion in *A. Litoralis* (Gramineae). *Physiology Plantarum* 47: 177-184.
- Popp M. 1984. Chemical composition of Australian mangroves. I. Inorganic ions and organic acids. *Zeitschr Pflanzenphysiologie* 113: 395-409.
- Popp M. 1994. Salt resistance in herbaceous halophytes and mangroves. In : Behnke HD, Luttge U & Esser K. (Eds.), *Progress in Botany*, Springer, Berlin, pp 416-429.
- Pujol JA, Calvo JF & Diaz LR. 2000. Recovery of germination from different osmotic conditions by four halophytes from southeastern Spain. *Annals of Botany* 85: 279-286.
- Rengel Z. 1992. The role of calcium in salt toxicity. *Plant Cell and Environment* 15: 625-632.
- Roth LC. 1992. Hurricane and mangrove vegetation: effects of Hurricane Joan, October 1988, on the vegetation of Isla del Venado-Bluefields, Nicaragua. *Biotropica* 24: 375-384.
- Schachtman DP, Kumar R, Schroeder JI & Marsh EL. 1997. Molecular and functional characterization of a novel low-affinity cation transporter (LCT1) in higher plants. *Proceedings of the National Academy of Sciences, USA*. 94: 11079-11084.
- Scholander PF. 1968. How mangroves desalinate seawater. *Plant Physiology* 21: 251-261.
- Shan L, RenChao Z, SuiSui D & SuHua S. 2008. Adaptation to salinity in mangroves: Implication on the evolution of salt tolerance. *Chinese Science Bulletin* 53: 1708-1715.
- Slater A, Scott N & Fowler M. 2003. *Plant Biotechnology. The genetic manipulation of plants*. Oxford University Press. New York.
- Stewart GR & Lee JA. 1974. The role of proline accumulation in halophytes. *Planta* 120: 279-289.
- Storey R, Ahmed N & Wyn Jones RG. 1977. Taxonomic and ecological aspects of the distribution of glycinebetaine and related compounds in plants. *Oecologia* 27: 319-332.
- Suarez N, Sobrado MA & Medina E. 1998. Salinity effects on the leaf water relations components and ion accumulation patterns in *Avicennia germinans* L. seedlings. *Oecologia* 114: 299-304.
- Tattini M, Gucci R, Coradeschi MA, Ponzio C & Everard JD. 1995. Growth, gas exchange and ion content in *Olea europaea* plants during salinity stress and subsequent relief. *Physiology Plant* 95: 203-210.
- Torres BC & Bingham FT. 1973. Salt tolerance of Mexican wheat. I. Effect of NO₃ and NaCl on mineral nutrition, growth and grain production of four wheats. *Soil Science Society of America Proceedings* 37: 711-715.
- Tozlu I, Moore GA & Guy CL. 2000. Effect of increasing

- NaCl concentration on stem elongation, dry mass production, and macro-and micro-nutrient accumulation in *Poncirus trifoliata*. *Australian Journal of Plant Physiology* 27: 35-42.
- Turner NC & Jones MM. 1993. Turgor maintenance by osmotic adjustment. In: Turner NC & Kramer PJ. (Eds.), *Adaptations of plants to water and high temperature stress. A Review and Evaluation* pp 87-103.
- Ungar IA. 1991. *Ecophysiology of vascular Halophytes*. Boca Raton, CRC. Press, Florida.
- Ungar IA. 1996. Effect of salinity on seed germination, growth and ion accumulation of *Atriplex patula* (Chenopodiaceae). *American Journal of Botany* 83: 604-607.
- Yeo AR. 1983. Salinity resistance: Physiologies and prices. *Physiology Plant* 58: 214-222.
- Zheng WJ, Wang WQ & Lin P. 1999. Dynamics of element contents during the development of hypocotyls and leaves of certain mangrove species. *Journal of Experimental Marine Biology and Ecology* 233: 247-257.