

COMUNICACIONES BREVES (Short notes)

***Satureja xdelpozoi*, nuevo híbrido del Sureste Peninsular**

Pedro Sánchez Gómez¹, Juan F. Jiménez² & Ramón Morales³

¹ Departamento de Biología Vegetal (Botánica), Universidad de Murcia, Campus de Espinardo, 30100, Murcia.

² Jardí Botanic, Universitat de Valencia, C/ Quart 80, 46008, Valencia.

³ Real Jardín Botánico de Madrid, Plaza de Murillo 2, 28014, Madrid.

Recibido: 1 Octubre 2005

Aceptado: 28 Octubre 2005

De acuerdo con la bibliografía consultada (López 1982), hasta el momento se conocen en la Península Ibérica 2 híbridos interespecíficos del género *Satureja* L. En este trabajo se describe un nuevo híbrido procedente del cruzamiento de *Satureja obovata* subsp. *obovata* y *Satureja intricata* subsp. *gracilis*, táxones que conviven en diversos puntos del Sureste Peninsular.

Descripción

Satureja xdelpozoi Sánchez Gómez, J.F. Jiménez & R. Morales **nothosp. nov.**

(*Satureja obovata* subsp. *obovata* x *S. intricata* subsp. *gracilis*)

Planta robusta, intermedia inter parentes. A Satureja obovata majoribus suis calicibus, versus 6 mm differt. A Satureja intricata corolla rosea differt.

Nothotaxon amico Enrique del Pozo Garnica *dicatur*

Habitat inter parentes in collibus calcareis, prope Letur (Albacete)

Holotypus. Albacete: Letur, Rambla de Sorbas, 30SWH7433, 1140 m, matorral sobre colubios dolomíticos, 08/10/2004, P. Sánchez Gómez & J.F. Jiménez, MUB 15736.

Planta de porte robusto, intermedia entre los parentales (fig 1). Presenta caracteres de ambos proge-

nitores. El porte es parecido a *Satureja intricata*, con verticilastros densos dispuestos en inflorescencias generalmente simples o escasamente ramificadas, aunque los tallos suelen presentar mayor longitud. Las flores presentan una corola rosada, carácter procedente de *Satureja obovata*, aunque con coloración más tenue. Los cálices presentan una longitud de hasta 6 mm, tamaño parecido a los de *S. intricata*, sin embargo, los dientes inferiores son claramente más cortos que la longitud del tubo, tal como ocurre en los de *S. obovata* con cálices tubular-campanulados. Las hojas presentan frecuentemente un ápice redondeado, como *S. obovata*, sin embargo pueden observarse algunas con dentición apical tal como ocurre en *S. intricata*. La pilosidad del carpostegio, presencia de glándulas, papilas y tonalidades moradas del cáliz, son caracteres próximos a *S. obovata*.

Agradecimientos

A D. Manuel López López por la diagnosis latina.

Referencias

López G. 1982. Compendio de las *Saturejæ* ibéricas con las noticias más notables referentes sobre todo a algunas de ellas. Anales del Jardín Botánico de Madrid 38 (2): 361-415.




Figura 1. *Satureja xdelpozoi*. a: hábito. b: cáliz, b1: *Satureja intricata* subsp. *gracilis*, b2: *Satureja xdelpozoi*, b3: *Satureja obovata* subsp. *obovata*.

Figure 1. *Satureja xdelpozoi*. a: habit. b: calyx, b1: *Satureja intricata* subsp. *gracilis*, b2: *Satureja xdelpozoi*, b3: *Satureja obovata* subsp. *obovata*.

Wild Siamese-twins in black tip sea catfish, *Arius dussumieri* (Valenciennes, 1840) from Gulf of Oman.

S M Al-Jufaily¹, Laith, A Jawad² & A N Al-Azi¹

(1) College of Agriculture and Marine Science, P.O. Box 34 Al-Khud 123, Sultanate of Oman. E-mail: sjufaily@squ.edu.om

(2) 15 Birkinshaw Grove, Riverstone Terraces, Upper Hutt, Wellington, New Zealand. (Corresponding author).

Received: 22 march 2005

Accepted: 7 july 2005

Siamese twins represent a condition in which two animals are conjoined to varying degrees. The occurrence of the phenomenon is well documented in different vertebrate groups (Witschi, 1952). Jussieu (1754) reported the first case among fish (*Carassius* sp.). It is common in both farmed and wild fish stock species (Hulata and Rothbard, 1978, Huang et al., 1987, Roberts, 1989, Bruno, 1990), where in the latter; only few cases survive beyond the absorbed yolk sac stage (Gordolla, 1968).

During a regular daily examination of the fishing yield in Al-Qareen village, Musnaa area, Batinah Region, 120 km north of Muscat, Gulf of Oman, Oman (June 2004), a twinned black tip sea catfish was found entangled in a traditional gillnet set at a depth of 3-5 m. They were in the post yolk sac-stage, fused against their caudal peduncle (Figure 1), and separate in an area in front of the anus. The twins measured 52.5 mm in total length. The nature of the fusion suggested that each of the two fish had separate internal organs except for a common anal opening. Each had its own dorsal and pectoral fins, and shared a common anal, pelvic and deformed caudal fins. The meristic characters of both twins are similar to those of the adult fish (Fischer & Bianchi, 1984).

A radiograph demonstrated the presence of overlapping caudal vertebral column for each fish (Figure 2). The two fish appeared to have separated from an area at the 21st vertebra (counting from the posterior end of the twins). Vertebral anomalies were noticed in the different body regions. The common caudal fin skeleton was completely deformed. All the hypurals and epurals bones were deformed and appeared in a single irregular bony structure. Vertebrae of the lower twin were compressed at the dividing region (Figure 2,3). The centra of vertebrae 1-4 and 13-15 of both fishes were deformed and fused together with their neural and haemal spines free and normal. The 8th vertebra (counting from the splitting

point) of the upper fish (Figure 4) had lost its posterior part and appeared compressed between the 7th and 9th vertebrae. Compressed, 11th & 12th vertebrae (counting from the dividing region of the lower fish (Figure 4) but not otherwise deformed were observed with their neural and haemal spines not deformed. Minor abnormalities including non-aligned and wavy neural and haemal spines of both fish from different body regions were also noted (Figures 2-4). The fin rays of the common anal fin were deformed and their connection to the underlying pterygiophores were lost. The last four fin rays of the dorsal fin of the upper fish was reduced in size while the remaining two appeared normal, but fused together. The dorsal fin rays of the lower fish were normal (Figures 2-4). The extracted otoliths of the upper fish are deeper than those of the lower fish.

Witsch (1952) suggested that over-ripeness of eggs leads to variety of developmental abnormalities resulted from decreased cohesion and interrelation between cells as well as between parts of inductive and of organ-forming regions. At this stage (post yolk-sac), there is no indication that the Siamese twin occurrence is a result of natural factors. On the other hand, the skeletal anomalies in the twins have various causes, but are usually considered to originate from mutations and from teratogenic effects of adverse environmental factors, such as mutagenic chemicals in the water (Longwell et al., 1992, Lien, 1997), or developing embryos and young individuals (Brown and Nunez, 1998, Vogel, 2000).

It seems likely that adverse environmental factors such as chemical pollution of the habitat may have played a role in producing skeletal abnormalities in the present twins. High levels of certain pollutants recorded from the Gulf of Oman (ROPME, 1999, De Mora et al., 2004) might have a direct effect on the developing stages. Pollution sources include thermal water from the power stations and hypersaline water from desalination plants (ROPME, 1999), wastewater




Figure 1. Conjoined twins of *Arius dussumieri* (Valenciennes, 1840). Scale bar =1cm




Figure 2. Radiograph of conjoined twins showing skeletal abnormalities. Scale bar = 1 cm.



Figure 3. Radiograph of conjoined twins showing skeletal abnormalities in the posterior region of the body. Scale bar =1cm.



Figure 4. Radiograph of conjoined twins showing skeletal abnormalities in the anterior region of the body. Scale bar = 1cm.

treatment sites (Al-Sabahi, 1997), solid waste (Linden et al., 1990), recreation and tourism facilities (Salm, 1989, 1991, 1993), and oil refineries (Al-Kharus, 1996). The main pollutants present in the coastal water of the Gulf of Oman are heavy metals and hydrocarbon contaminants and organic pollutants (Fowler, 2002) such as the pesticide DDT, which has been implicated as a cause of skeletal deformities in other fish species (Bengtson et al., 1985).

Vertebral deformities such as those reported here in the Siamese twin may affect the biology of the fish indirectly through inhibiting its free-swimming movements (Sadler, 1990). An increased incidence of skeletal deformities among commercial fish species would suggest environmental deterioration and hence signal the need for prompt remedial action.

Acknowledgements

We would like to thank Bruce Marshall and Raymond Coory, both of the Museum of New Zealand Te Papa Tongarewa, respectively for continuing advice and suggestions in the preparation of the manuscript and for technical help in preparing the figure.

References

- Al-Kharusi L.H.S. 1996. Distribution and accumulation of oil pollutants on the region of Mina Al-Fahal in the Sultanate of Oman and their effects on the marine environment. M.Sc. thesis, Suez Canal University, Egypt, 136p.
- Al-Sabahi A.M. 1997. Wastewater treatment and reuse in the Sultanate of Oman. Third Gulf Water Conference, Muscat, March 1997, Vol. 3, p.859-888.
- Bengtson BE, Bengtson A & HinbergM. 1985. Fish deformities and pollution in some Swedish waters. *Ambio* 14: 32-35.
- Brown CL & Nunez J.M. 1998. Disorders of development. In *Fish diseases and disorders*. Vol. 2 (Leatherland JE & Woo PTK, eds) Oxon: CABI Publishing, p. 340.
- Bruno DW. 1990. Occurrence of a conjoined twin among farmed Atlantic salmon, *Salmo salar* L. *Journal of Fish Biology* 37: 501-502.

- De Mora SJ, Fowler SW, Wyse E & Azemard S. 2004. Distribution of heavy metals in marine bivalves, fish and coastal sediments in the Gulf and Gulf of Oman. *Marine Pollution Bulletin* 49: 410-424.
- Fischer W & Bianchi G. 1984. FAO species identification sheets for fishery purposes. Western Indian Ocean: Fishing area 51. FAO, 6 volumes.
- Fowler SW. 2002. Agrochemicals. In *The Gulf ecosystem: health and sustainability* (Khan NY, Munawar M & Price ARG). Bakhuy Publishers, Leiden, p.193-217.
- Gordolla PH. 1968. Fortpflanzungsfähige siamesische Zwillinge bei *Poecilia reticulata*. *Monatsschrift fuer Ornithologie und Vivarienkunde Ausgabe B., Aquarien Terrarien* 15: 407-409.
- Huang CM, Cheng HJ, Chang SL, Hao NH & Liao IC. 1987. Siamese twins in tilapia. *Journal of Fish Biology* 31: 441-442.
- Hulata G & Rothbard S. 1978. 'Siamese-twins' in *Sarotherodon mossambicus*. *Journal of fish Biology* 13: 45.
- Jussieu LC 1754. Observation sur deux petit poissons reunis par le ventre. *Memoires de l'Academie Sciences de Paris* 1754: 30.
- Lien NTH 1997. morphological abnormalities in African catfish *Clarius gariepinus* larvae exposed to malathion. *Chemosphere* 35 1475-1486.
- Linden O, Abdulraheem MY, Gerges MA, Alam L, Behbehani M, Borhan MA & Al-Kassab F. 1990. State of the marine environment in the ROPME sea area. UNEP Regional Seas Reports and Studies No. 112, Rev. 1, UNEP.
- Longwell AC, Chang S, Hebert A, Hughes JB & Perry D. 1992. Pollution and developmental abnormalities of Atlantic fishes. *Environmental Biology of Fishes* 35: 1-21.
- Roberts RJ. 1989. *Fish Pathology*. 2nd edition. London: Bailliere Tindall.
- ROPME, 1999. Overview on land-based sources and activities affecting the marine environment in the ROPME sea area. UNEP Regional Seas Reports and Studies No. 168.
- Sadler TW. 1990. *Langman's medical embryology*. 6th edition. Baltimore: Williams & Wilkins, p. 411.
- Salm RV. 1989. A general overview of the coast between Ra's al Hadd and Dhofar. IUCN Coastal Zone Management Project 9429. Sultanate of Oman, CZMP3: F6:3p.
- Salm RV. 1991. Oman coastal zone management program: A framework for action. IUCN, Gland, Switzerland.
- Salm RV. 1993. Coral reefs of the Sultante of Oman. *Atoll Research Bulletin* 380: 1-85.
- Vogel G. 2000. Zebrafish earns its strips in genetic screens. *Science* 288: 1160-1161.
- Witschi, E. 1952. Over-ripeness of the eggs as a cause of twinning and teratogenesis: a review. *Cancer Research* 12: 763-786.

